



26GT[®]

Product information sheet

- // Active Ingredients: Iprodione (23.3%); equivalent to 2 lb. iprodione per gallon
- // FRAC Code: 2 (Dicarboximide)
- // Systemicity: Localized penetrant
- // Formulation: Oil in water emulsion
- // Signal Word: Caution

26GT makes a great rotational partner with Exteris[®] Stressgard[®] to help manage resistance to SDHI chemistry (for more information on resistance mitigation visit FRAC website <http://www.frac.info/resistance-overview>)

Best Uses

26GT[®] provides control of a wide range of fungal diseases including dollar spot, brown patch, snow molds, Microdochium patch, leaf spots, Curvularia blight

Key Strengths

Control of DMI-resistant dollar spot, summer safety, leaf spot on bermudagrass

Attributes of 26GT

- // Year-round, broad-spectrum disease control
- // Non-DMI chemistry avoids plant growth regulator (PGR) effects
- // Easily pairs with PGR programs for flexible use
- // Advanced formulation for quick dollar spot knockdown

Solutions for Tough Diseases

- // 26GT controls DMI- and SDHI- resistant dollar spot
- // 26GT can be used for summer dollar spot and brown patch control without negative turf growth effects
- // Combine 26GT with Signature[™] XTRA Stressgard or Chipco[®] Signature for summer decline management
- // 26GT controls spring and fall diseases such as leaf spot and Microdochium patch
- // Use 26GT for pink and gray snow mold control

How to Use 26GT


Apply 26GT as part of an integrated management program on greens, tees and fairways. Apply preventatively when conditions are favorable for disease. 26GT, Signature™ XTRA™ Stressgard and Chipco® Signature can be tank-mixed for management of summer decline. For additional disease control and plant health effects, use products containing Stressgard such as Interface® Stressgard.

Key Diseases Controlled ¹	Rate (per 1,000 sq ft)	Application Interval
Dollar Spot ••	3.0 - 4.0 fl oz 2.0 - 4.0 fl oz	14 - 21 days (greens & tees) 14 - 28 days (fairways & other turf areas)
Brown Patch • Bipolaris and Dreschlera Leaf Spots ••	3.0 - 4.0 fl oz	14 - 21 days (greens & tees) 14 - 28 days (fairways & other turf areas)
Red Thread • Curvularia Blight ••	4.0 fl oz 4.0 - 8.0 fl oz (Curvularia control in bermudagrass only)	14 days
Microdochium (Fusarium) Patch ••	4.0 - 8.0 fl oz	14 - 21 days
Snow Molds •	4.0 - 8.0 fl oz	One application before winter snow cover; if possible, another application during a mid-winter thaw
Summer Decline ••	3.0 - 4.0 fl oz 26GT® + 4.0 - 8.0 oz Chipco® Signature™ or 2.0 - 6.0 oz Signature XTRA Stressgard	14 days
Large Patch •	4.0 fl oz	14 - 21 days

• = cool-season turf diseases

•• = warm-season turf diseases

¹See the 26GT label for the complete list of diseases controlled. Refer to the 26GT label for application information. Do not apply more than 35.0 fl. oz./1,000 sq. ft. Do not exceed 6 applications per year. Always read and follow label instructions carefully.



University of Connecticut research showing rainfastness of 26GT. 2.0 fl oz of 26GT applied and followed up with 0.10 inches of rain 30 minutes post-application. (Bayer)

www.EnvironmentalScience.Bayer.US | [@BayerGolf](https://twitter.com/BayerGolf)



ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS
Bayer Environmental Science, a Division of Bayer CropScience LP, 5000 CentreGreen Way, Suite 400, Cary, NC 27513. For additional product information, call toll-free 1-800-331-2867. Not all products are registered in all states. Bayer (reg'd), the Bayer Cross (reg'd), Chipco®, 26GT®, Exteris®, Stressgard®, Interface®, Chipco® and Signature™ are trademarks of Bayer. © 2019 Bayer CropScience LP. ES-1018-26GT-427-A-R1