

<p>PRECAUTIONARY STATEMENTS</p> <p>HAZARDS TO HUMANS AND DOMESTIC ANIMALS</p> <p>“Do not apply granular lawn products by hand. Avoid contact with hands or skin.”</p> <p>“Broadcast applications must NOT be made using hand-held devices such as a belly grinder or handheld rotary applicator. Such equipment may only be used for spot treatments” “ Maximum rate application turfgrass [including lawns] is [4 lbs per 1,000 sq.ft.—that reflects an maximum application rate of 2 pound active ingredient per acre) “Maximum of two applications per year.)</p> <p>CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eves, skin, or clothing. Prolonged or frequently repeated skin contact cause allergic reactions in some individuals.”</p>	
FIRST AID	
If Swallowed:	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have a person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person</li></ul>
If in eyes:	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li><li>• Remove contact lenses, if present, after 5 minutes, then continue rinsing.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
If on skin or clothing:	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call poison control center or doctor for treatment advice.</li></ul>
If inhaled:	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 91 1 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements.	
<p>ENVIRONMENTAL HAZARDS</p> <p>Atrazine can travel (seep or leach) through soil and can enter ground water, which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very penneable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.</p> <p>This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning of equipment or disposing of equipment washwaters.</p>	

NOTICE

Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. To the extent consistent with applicable law, Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. To the extent consistent Jaw, In no use shall Sunniland Corporation ml liable for consequential, special, or indirect damages resulting from the use or handling of this product

Manufactured by:

SUNNILAND CORPORATION

P.O. BOX 8001

SANFORD, FL 32772-8001

EPA REG. NO. 9404-82

EPA EST. NO. 9404-FL-1

# SUNNILAND WEED & FEED

## For St. Augustine lawns 25-4-6

### BROADLEAF AND GRASSY WEED CONTROL WITH PREMIUM LAWN FOOD

20 Pounds of WEED & FEED covers 5,000 sq.ft. of lawn

Prevents Crabgrass, Paspalums and Spotted Spurge

Controls Florida Pusley, Lippia, Pigweed and as many as 44 other Broadleaf and/or Grassy Weeds as listed.

## FOR USE ON ST. AUGUSTINE AND CENTIPEDE LAWNS

GUARANTEED ANALYSIS	F99	094972
Total Nitrogen (N)*.....		25.00%
8.30% Ammoniacal Nitrogen		
16.70% Urea Nitrogen		
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ).....		4.00%
Soluble Potash (K <sub>2</sub> O).....		6.00%
Chlorine, Not more than .....		6.00%

\* Contains 12.52 units of slow release nitrogen.

Derived From: Poly Sulphur Coated Urea, Ammonium Sulphate, Diannnonium Phosphate, Muriate of Potash.

ACTIVE JNGREDIENT:	
Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine .....	1.15%
Related compounds .....	0.00%
INERT INGREDIENTS:.....	98.85%
TOTAL.....	100.00%

This product contains 22 lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lbs. Atrazine per acre when used according to label.

KEEP OUT OF REACH OF CHILDREN

CAUTION

NET WEIGHT 50 LB (22.68 kg)

## DIRECTIONS FOR USE

It is a Violation of Federal law to use this product in a manner inconsistent with its labeling Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. This product will control both emerged weeds and weeds from seeds. Some moisture is required to activate the herbicide and the fertilizer. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people, pets out of area during application.

## WEEDS CONTROLLED OR SUPPRESSED

Annual Bluegrass (Poa Annual)	Crowfoot Grass	Paspalums
Annual Sowthistle	Cutleaf Evening Primrose	Red Sorrel
Clover	Dichondra	Pigweed
Black Medic (yellow trefoil)	Dollarweed	Purslane
Blue Toadflax	Florida Belony (Rattlesnake Weed)	Rescuegrass
Bull Paspalum	P.orida Beggarweed Florida Pusley	Sandspur
Buckhorn Plantain	Fragrant Cudweed (Rabbit Tobacco)	Shepherdspurse
Bur Clover	Henbit	Sickelpod
Carolina Geranium (Cranesbill)	Knotweed	Spotted Spurge
Chickweed (Common & Mouse-Ear)	Lespedeza	Spurweed
Cockebur	Mat Lippia (Matchweed)	Stinging Nettle
Common Dandelion	Moneywort	Swinecrest
Common Purslane	Morning Glory	Venus Looking Glass
Grabgrass	Narrowleaf Vetch	Virginia Creeper
Cudweed	Oxalis (Common and Yellow Woodsorrel)	Wild Mustard
		WoodSorrel

## WHEN TO APPLY

Best weed control will result if applied before weeds sprout since the product acts as a pre-emergent and also a post-emergent weed killer. Established weeds are harder to kill and may require an additional application in the fall or spring. Apply January 1 and March 1 for the best control otherwise product may be applied up to June 15 with good weed Control. A fall application will be needed for control certain annuals such as Henbit, Chickweed, Rabbit Tobacco, Florida Betony and Evening Primrose. In case of heavy weed infestation or low soil moisture a repeated application may be necessary. Kill time of mature weeds may be 4 to 5 weeks. Wait 2 months for repeat application. Apply when air is calm to prevent particle drift. Do not make more than 2 applications per year.

## HOW TO APPLY

This product maybe applied with either a drop type or broadcast type spreader. Apply evenly across the entire lawn at a rate of 20 pounds to treat 5,000 sq.ft. (equivalent to 4 lbs per 1,000 sq.ft.) Do not exceed recommended rates. Apply to dry grass. Sprinkle the area immediately after application to wash granules off the grass blades down to soil surface.

## Spreader Settings - 20lbs / 5,000 sq.ft

Cyclone 2.75 Scott 5.5

Note: spreader rate varies with walking speed.

Check spreader over known area.

## WHERE AND WHEN TO USE

This material Is for use on St. Augustine and Centipede Lawns only. On newly seeded centipede, wait until grass has been mowed at least once. Do not plant sprigs within one month after application, or seed within 6 months after application. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry.

DO NOT USE ON BLUEGRASS, FESCUE, BENTGRASS, BAHIA RYE BERMUDA, OR DICHONDRA LAWNS.

## USE PRECAUTIONS

Do not apply this product on or over the root zone of ornamental trees and shrubs, nor to bedding plants, flowers, or garden plants since injury to these plants may result. Do not apply to lawns containing grasses or mixtures of grasses other than St. Augustine or Centipede grasses as recommended on this label or injury to those other grasses may result. Do not enter or allow others to enter treated areas, (except those involved with the watering),until the watering-in is complete and the surface is dried. DO NOT APPLY BY HAND. Do not allow product to come in contact with hands. Application of this product by hand-held devices (e.g. belly grinder) is limited to spot treatment Wear long-sleeved shirt, long pants, shoes and socks when applying this product. Restrictions: Do not apply this product in a way that will contact any person or pet either directly or through drift. Keep people and pets out of area during application. \*This product must be watered in immediately after application. \*\*Entry Restrictions: Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry.”

## STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, food stuffs, feed, or seed by storage or disposal. PESTICIDE DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions. CONTAINER DISPOSAL IF EMPTY: Do not reuse this bag. Place in trash or offer for recycling if available. IF PARTLY FILLED: Call your local waste agency or 1-800- CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. SPILL: In case of spill, sweep up and dispose of material according to PRODUCT DISPOSAL directions listed above.