

GARY'S GREEN®

18-3-4

GRIGG Gary's Green is the flagship of GRIGG's Proven Foliar Nutrient line – offering the ultimate in quality, performance and compatibility. It contains three sources of nitrogen, potassium, phosphorus and a micronutrient package. It improves turf color response, density and vigor. It is widely used as a primary source of N for effective 'spoon feeding' in turf nutrition programs.

Also available in a phosphate free version.

Key Advantages

- Contains soluble nutrients for efficient uptake and use
- Proprietary formulation provides fast nitrogen uptake;
 N promotes consistent turfgrass shoot growth
- Phosphorus plays a role in plant metabolic processes that transfer energy throughout the plant
- Potassium regulates primary physiological processes that impact turf response to stress and supports cellular processes that impact photosynthesis, water regulation, respiration and protein production
- Recommended for golf and sports turf

Application and Use

Cool Season Grasses: Apply 6-9 fl oz per 1,000 sq ft or 2-3 gal per acre [20-30 L per hectare] as needed every 7-14 days.

Warm Season Grasses: Apply 9-12 fl oz per 1,000 sq ft or 3-4 gal per acre [30-40 L per hectare] as needed every 7-14 days.

For a distributor near you contact: 800 300 6559 or www.grigg.co

GRIGG is part of Brandt Consolidated, Inc. 2935 South Koke Mill Road Springfield, IL 62711 www.brandt.co

Guaranteed Analysis
Total Nitrogen (N)
2.00% Ammoniacal nitrogen
1.50% Nitrate nitrogen
14.50% Urea nitrogen
Available Phosphate (P ₂ O ₅)
Soluble Potash (K ₂ O)
Magnesium (Mg)
0.50% Water soluble magnesium
Copper (Cu)
0.12% Water soluble copper
Iron (Fe)
1.00% Water soluble iron
Manganese (Mn)0.10%
0.10% Water soluble manganese
Zinc (Zn)0.10%
0.10% Water soluble zinc

Derived from urea ammonium nitrate, monoammonium phosphate, urea, potassium citrate, magnesium amino acid complex, copper amino acid complex, iron amino acid complex, manganese amino acid complex and zinc amino acid complex.

Make frequent applications at lower rates, or apply higher rates at times of greater plant demand. Optimum rate of application will vary depending on treatment interval, soil properties (such as pH, organic matter content, texture), weather conditions, time of year, plant species and its nutrient requirements. For best results, follow soil/tissue test recommendation.

