



RightLine®

PROHEX 27.5 WDG

Plant growth regulator
for use on turf*

*Not for use in California

ACTIVE INGREDIENT:

Prohexadione-calcium**

[calcium 3-oxido-5-oxo-4-propionylcyclohex-3-enecarboxylate].....27.5%

Other Ingredients:.....72.5%

Total:.....100.0%

** Equivalent to 0.275 pounds of active ingredient per pound of product

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Poison Control Hotline at 1-800-222-1222 for emergency medical treatment information 24 hours a day, seven days a week.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container only. After partial use, close the container tightly. Store in a secure place that is cool and dry. Use spray and stock solutions within 24 hours. Immediate use is required if another component is added to the spray solution.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, if available, or puncture or dispose of in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

See inside panel for additional Precautionary Statements

EPA Reg. No. 62097-41-93051 Est. No. 39578-TX-001

Distributed By:

Rightline LLC
950 Falcon Drive
Malden, MO 63863
Tel: 770-335-3015

NET CONTENTS: 6 LBS.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves made of any waterproof material polyethylene or polyvinyl chloride

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE items separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement, specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment (PPE) and restricted entry intervals. The requirements in this box only apply to uses of this product covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval level (REI) of 12 hours unless wearing appropriate PPE.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, plants, soil or water is:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical resistant gloves made of any waterproof material

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep unprotected persons out of treated areas until sprays have dried.

TURF (all approved states except California):

Rightline PROHEX 27.5 WDG can be used as a production management tool on turf to reduce vegetative growth. The active ingredient in Rightline PROHEX 27.5 WDG, prohexadione-ca, reduces the growth of turf by inhibiting the biosynthesis of gibberellin, a natural plant hormone that regulates cell elongation. The inhibition of gibberellin slows the growth of turf which will result in reduced frequency of mowing and amounts of grass clippings in treated areas. Following an application of Rightline PROHEX 27.5 WDG to turf, a reduction in growth and grass clippings will typically be observed by three days.

Rightline PROHEX 27.5 WDG can be applied to turf that is green and actively growing. Under conditions of stress, applications need to be delayed. Apply lower rates if the turf

is entering stressful growing conditions like pest pressure, high temperatures, low moisture conditions or certain cultural practices including, but not limited to, aeration or verticutting. Under certain environmental conditions, including those that may occur early in the season, transient bronzing may be observed. For information on application scheduling, rates and timings on turf for Rightline PROHEX 27.5 WDG see "Table 1: Application Rates & Timings for Using a Set/Calendar Spray Schedule" and "Table 2: Application Rates and Timings Using Growing Degree Days (GDD) to Determine Spray Timings" for Rightline PROHEX 27.5 WDG in Turf located below.

For consistent performance on turf, add one pound of a spray grade ammonium sulfate (AMS) for every pound of Rightline PROHEX 27.5 WDG applied. Use a standard spray adjuvant, preferably a non-ionic surfactant, to improve leaf coverage and performance consistency. Follow the manufacturer's rate directions.

Mandatory Spray Drift

Aerial Applications

- Do not release spray at a height greater than 10 ft above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For all applications, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Ground Applications

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Spray Drift Advisories

- THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
- BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

• IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

- Adjust Nozzles - Follow nozzle manufacturer's recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.
- BOOM HEIGHT - Ground Boom
Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the turf and have minimal bounce.
- RELEASE HEIGHT - Aircraft
Higher release heights increase the potential for spray drift. When applying aurally to target area, do not release spray at a height greater than 10 feet above the target area, unless a greater application height is necessary for pilot safety.
- SHIELDED SPRAYERS
Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area. TEMPERATURE AND HUMIDITY When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.
- TEMPERATURE INVERSIONS
Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally

in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

• **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Ground Application: Apply Rightline PROHEX 27.5 WDG to actively growing turf using calibrated application equipment. Rightline PROHEX 27.5 WDG is taken up through the leaves of the plants. So, good spray coverage of the plant foliage is essential to get good uptake and consistent growth suppression. To achieve good coverage, apply Rightline PROHEX 27.5 WDG in a sufficient spray volume utilizing proper spray pressure, nozzles, nozzle spacing and applicator speed. Apply Rightline PROHEX 27.5 WDG in a spray volume of 1 to 2 gallons water per 1000 sq. ft. but it can be applied in a spray volume of 0.5 to 5.0 gallons water per 1000 sq. ft. Avoid applying Rightline PROHEX 27.5 WDG during warm, low humidity or windy conditions. The absorption of the spray mix is improved with longer drying times.

Maintain constant agitation during mixing of the spray solution and application. Do not allow the spray tank to sit for prolonged periods without any agitation. Do not allow the spray mix to sit in the tank overnight.

Mowing & Irrigation: Mowing and irrigation events can affect the efficacy of the Rightline PROHEX 27.5 WDG applications. After mowing, do not make an application of Rightline PROHEX 27.5 WDG for at least one hour. Following an application of Rightline PROHEX 27.5 WDG, do not mow the treated turf for at least one day and do **not** apply irrigation for 4 hours.

Turf Maintenance: Following applications of Rightline PROHEX 27.5 WDG, maintain the areas of turf that were treated as normal with proper levels of irrigation and fertility. Implement a good pest control plan as necessary to maintain the turf in a healthy condition. If tank mixing with other product(s) precaution is advised. The user needs to test tank mixes that are unfamiliar on a small area before the treatment is applied to larger areas. Test tank mixes for compatibility before mixing a larger quantity for an application. User assumes all risk for tank mixes as it is not possible for manufacturer to test all possible tank mix combinations for compatibility, safety and effect on efficacy of the products. It is the pesticide user's responsibility to

ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations, and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Application Timing Methods

Rightline PROHEX 27.5 WDG may be applied to control vegetative growth in turf by two different application timing methods in turf. The product may be applied to the turf on a set/calendar spray schedule at regular intervals or it may be applied by assessing potential for turf growth using **Growing Degree Days (GDD)** to time the applications to control turf growth. These methods allow the person(s) managing the turf to select the best and most efficient use method of Rightline PROHEX 27.5 WDG for their particular facility.

Rightline PROHEX 27.5 WDG may be applied on a regular set/calendar schedule to maintain an even turf regulation or height control. Make the first application when turf is green and actively growing. Instructions for making repeat applications are provided in Table 1 below by turf type, species and growing area (Fairways, Roughs, Greens and/or Tees).

When applying Rightline PROHEX 27.5 WDG using the Growing Degree Day (GDD) model, use the environmental data from the location or immediate area. Make the first application of Rightline PROHEX 27.5 WDG once the turf is green and actively growing. Research has shown that 280-350 GDD needs to be used for re-application timings for common golf situations to assure even turf regulation and to avoid rebound growth. Table 2 below details the application rates and timings when using GDD to determine application timings.

With either application scheduling method, delay applications to the turf when the turf is under conditions of stress. Apply lower rates if the turf is entering stressful growing conditions including pest pressure, high temperatures, low moisture conditions or certain cultural practices including, but not limited to, aeration or verticutting. Under certain environmental conditions, including those that may occur early in the season, transient bronzing may be observed.

Table 1. Application Rates & Timings for Using a Set/Calendar Spray Schedule

	Rightline PROHEX 27.5 WDG Applications to Slow Vertical Growth, Reduce Mowing, Improve Color & Quality			
	Fairway & Rough Areas (Application Intervals: 2-4 Weeks)		Green & Tee Areas (Application Intervals: 1-2 Weeks)	
	oz/acre	oz/1000 sq. ft.	oz/acre	oz/1000 sq. ft.
Warm Season Turf*				
Hybrid Bermudagrass including Champion, MiniVerde, Riviera, TifEagle, TifGreen, TifSport, Tifway & 419	29.1 – 43.6 (0.50-0.75 lbs ai)	0.67 – 1.0 (0.013-0.019 lbs ai)	7.25 – 14.5 (0.13-0.25 lbs ai)	0.17 – 0.34 (0.003-0.006 lbs ai)
Zoysiagrass cultivars including Empire, Zenith, Emerald, El Toro and Meyer St. Augustine spp. Paspalum spp. Kikuyugrass spp.	29.1 – 43.6 (0.50-0.75 lbs ai)	0.67 – 1.0 (0.013-0.019 lbs ai)		
Cool Season Turf				
Bentgrass: L-93, Penn A4, Penn G2, Pennlinks, Penneagle, Southshore, Declaration, Independence and SR series	7.25 – 14.5 (0.13-0.25 lbs ai)	0.17 – 0.34 (0.003-0.006 lbs ai)	1.8 – 7.25 (0.03-0.13 lbs ai)	0.04 – 0.17 (0.0006-0.003 lbs ai)
Annual & Perennial Poa annua cultivars	5.43 – 10.87 (0.09-0.19 lbs ai)	0.125 – 0.25 (0.002-0.004 lbs ai)	1.8 – 7.25 (0.03-0.13 lbs ai)	
Kentucky bluegrass: Midnight, Odyssey	14.5 – 21.8 (0.25-0.38 lbs ai)	0.34 – 0.50 (0.006-0.009 lbs ai)		
Perennial ryegrass: Manhattan IV Tall Fescues Fine Fescues	21.8 – 29.1 (0.38-0.50 lbs ai)	0.50 – 0.67 (0.009-0.01 lbs ai)		

*Lower rates may be applied for color and quality enhancement. When using shorter application intervals, evaluate performance using the lower directed rates.

Table 2. Application Rates & Timings Using Growing Degree Days (GDD) to Determine Spray Timings.

	Rightline PROHEX 27.5 WDG Applications to Slow Vertical Growth, Reduce Mowing, Improve Color & Quality			
	Fairway & Rough Areas (280-350 GDD Interval)		Green & Tee Areas (280-300 GDD Interval)	
	oz/acre	oz/1000 sq. ft.	oz/acre	oz/1000 sq. ft.
Warm Season Turf*				
Hybrid Bermudagrass including Champion, MiniVerde, Riviera, TifEagle, TifGreen, TifSport, Tifway & 419	12 – 24 (0.21-0.41 lbs ai)	0.28 – 0.55 (0.005-0.009 lbs ai)	8 – 16 (0.14-0.28 lbs ai)	0.18 – 0.37 (0.003-0.006 lbs ai)
Zoysiagrass cultivars including Empire, Zenith, Emerald, El Toro and Meyer St. Augustine spp. Paspalum spp. Kikuyugrass spp.	12 – 24 (0.21-0.41 lbs ai)	0.28 – 0.55 (0.005-0.009 lbs ai)		
Cool Season Turf				
Bentgrass: L-93, Penn A4, Penn G2, Pennlinks, Penneagle, Southshore, Declaration, Independence and SR series	6 – 12 (0.11-0.21 lbs ai)	0.14 – 0.28 (0.003-0.005 lbs ai)	2 – 8 (0.03-0.14 lbs ai)	0.05 – 0.18 (0.0006-0.003 lbs ai)
Annual & Perennial Poa annua cultivars	4 – 8 (0.07-0.14 lbs ai)	0.09 – 0.18 (0.001-0.003 lbs ai)	2 – 8 (0.03-0.14 lbs ai)	
Kentucky bluegrass: Midnight, Odyssey	8 – 16 (0.14-0.28 lbs ai)	0.18 – 0.37 (0.003-0.006 lbs ai)		
Perennial ryegrass: Manhattan IV Tall Fescues Fine Fescues	8 – 16 (0.14-0.28 lbs ai)	0.18 – 0.37 (0.003-0.006 lbs ai)		

*Lower rates may be applied for color and quality enhancement. When using shorter application intervals, evaluate performance using the lower directed rates.

Table 3. Applications for Overseeding Enhancement in Bermudagrass.

Turf Type	oz/acre	Application Interval	Notes
Hybrid Bermudagrass including Champion, MiniVerde, Riviera, TifEagle, TifGreen, TifSport, Tifway, Yukon & 419	1 – 2 (0.02-0.03 lbs ai)	Apply one application of Rightline PROHEX 27.5 WDG 3-5 days prior to seeding. Any preparation of the site including verticutting, spiking or scalping needs to be done 1-2 days after the application.	Rightline PROHEX 27.5 WDG is a foliar- active product. Therefore, germination and seedling growth of the over-seeding are not affected when used as directed. Use good cultural practices, including fertilization and irrigation, in conjunction with Rightline PROHEX 27.5 WDG to optimize results. Some temporary yellowing of the bermudagrass turf may be observed, however, this will not affect the over- seeding.

Table 4. Applications for Reduction of Poa annua in Bermudagrass Overseeding.

Turf Type	oz/acre	Application Interval
Hybrid Bermudagrass (including Tifway 419, TifSport & Riviera	0.90 – 1.75 (0.02-0.03 lbs ai)	Reapply Rightline PROHEX 27.5 WDG at 3-4 week intervals during periods of active Poa annua growth.

Limitation on Turf:

- Rightline PROHEX 27.5 WDG is rainfast 1 hour after application.

Note: If calcium or boron are tank mixed with Rightline PROHEX 27.5 WDG, the resulting growth control may be less than favorable.

Restrictions on Turf:

- **DO NOT** apply more than 1132.6 oz of Rightline PROHEX 27.5 WDG (19.47 lbs ai) per acre per year.
- **DO NOT** apply more than 26 oz of Rightline PROHEX 27.5 WDG (0.45 lbs ai) per 1000 sq. ft. per year.
- **DO NOT** apply more than 1 oz per 1000 sq. ft. or 43.6 oz per acre (0.02 lbs ai) per 1000 sq. ft. or 0.75 lbs ai per acre of Rightline PROHEX 27.5 WDG in a single application.
- **DO NOT** retreat in less than 1 week.
- **DO NOT** make more than 52 applications per year when applied at lower rate.
- **DO NOT** apply Rightline PROHEX 27.5 WDG to turf that has been sodded or sprigged until it has knitted down and rooted firmly.
- **DO NOT** apply to turf growing in conditions that are not ideal for good growth. Also, DO NOT apply to turf that shows injury (leaf phytotoxicity) produced by any other prior pesticide applications as this type of injury can be enhanced or prolonged. Refer to Tank Mixing Information for additional tank mixing instructions and precautions.
- **DO NOT** graze treated areas or feed clippings previously treated with Rightline PROHEX 27.5 WDG to livestock.
- **DO NOT** apply Rightline PROHEX 27.5 WDG to turf using any type of irrigation/chemigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container only. After partial use, close the container tightly. Store in a secure place that is cool and dry. Use spray and stock solutions within 24 hours. Immediate use is required if another component is added to the spray solution.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, if available, or puncture or dispose of in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of RightLine LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold RightLine LLC and Seller harmless for any claims relating to such factors.

RightLine LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product

contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or RightLine LLC, and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, RIGHTLINE LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither RightLine LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF RIGHTLINE LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF RIGHTLINE LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

RightLine LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of RightLine LLC.



RightLine®

PROHEX 27.5 WDG

Plant growth regulator
for use on turf*

*Not for use in California

ACTIVE INGREDIENT:

Prohexadione-calcium**

[calcium 3-oxido-5-oxo-4-propionylcyclohex-3-enecarboxylate].....27.5%

Other Ingredients:.....72.5%

Total:.....100.0%

** Equivalent to 0.275 pounds of active ingredient per pound of product

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Poison Control Hotline at 1-800-222-1222 for emergency medical treatment information 24 hours a day, seven days a week.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container only. After partial use, close the container tightly. Store in a secure place that is cool and dry. Use spray and stock solutions within 24 hours. Immediate use is required if another component is added to the spray solution.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, if available, or puncture or dispose of in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

See inside panel for additional Precautionary Statements

EPA Reg. No. 62097-41-93051 Est. No. 39578-TX-001

Distributed By:

Rightline LLC
950 Falcon Drive
Malden, MO 63863
Tel: 770-335-3015

NET CONTENTS: 6 LBS.

PEEL BACK BOOK HERE