

<b>PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS</b> “Maximum rate per application turfgrass (including lawns) is 5lbs. Per 1,000 sq.ft. that reflects a maximum application rate of 2 pound active ingredient per acre] Maximum of two applications per year.”	
<b>CAUTION:</b> Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing!	
<b>FIRST AID</b>	
<b>If Swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have a person sip a glass of water if able to swallow .</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes .</li> <li>• Call poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements.	
<b>ENVIRONMENTAL HAZARDS</b>	
Atrazine can travel (seep or leach) through soil and can enter ground water, which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very penneable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water. This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply on or under ornamental trees, shrubs, budding plants, flowers or garden plants since injury may result.	

**NOTICE**

Sunniland Corporation warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. To the extent consistent with applicable law, Sunniland Corporation makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. To the extent consistent law, In no use shall Sunniland Corporation ml liable for consequential, special, or indirect damages resulting from the use or handling of this product

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turfgrass being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds

**STORAGE AND DISPOSAL**

\*Do not contaminate water, foodstuffs, feed or seed by storage or disposal\*

**PESTICIDE STORAGE:** Store this product in its original container in a dry, cool, secured area. **PESTICIDE PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions. **IF EMPTY:** Do not reuse this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800- CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **SPILL:** In case of spill, sweep up and dispose of material according to **PRODUCT DISPOSAL** directions listed above.

Manufactured by:  
**SUNNILAND CORPORATION**  
**P.O. BOX 8001**  
**SANFORD, FL 32772-8001**

**EPA REG. NO. 9404-51**

**EPA EST. NO. 9404-FL-1**

# SUNNILAND WEED & FEED

## 16-0-8

### For St. Augustine, Centipede & Zoysia Grass

- 25 Pounds WEEDS & FEEDS 5,000 sq. ft.
- For St. Augustine, Centipede, and Zoysia Grass
- WEEDS as it FEEDS. May be applied twice/year
- ALSO PREVENTS CRABGRASS GERMINATION

**Item # 094692**

### USE ON ST. AUGUSTINE, CARPETGRASS, CENTIPEDE AND ZOYSIA LAWNS FOR CONTROL OF WEEDS AS LISTED ON SIDE PANEL

<b>GUARANTEED ANALYSIS*</b>	<b>F99</b>
Total Nitrogen (N).....	16.00%
5.00% Ammoniacal Nitrogen	
1 1.00% Urea Nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>3</sub> ).....	0.00%
Soluble Potash (K <sub>2</sub> O).....	8.00%
Total Magnesium as Mg .....	1.20%
1.20% Water Soluble Magnesium as Mg	
Boron as B.....	0.02%
Chlorine, (Cl) Not more than.....	6.00%
Total Copper as Cu .....	0.05%
Total Iron as (Fe).....	3.00%
Total Manganese as (Mn).....	0.06%
Molybdenum as Mo .....	0.0005%
Total Zinc as Zn .....	0.05%
Derived From: Poly Sulphur Coated Urea, Ammonium Sulphate, Muriate of Potash, Sulphate of Potash Magnesia, Sodium Borate, Copper Oxide, Manganese Oxide, Iron oxide, Zinc Oxide, Sodium Molybdate.	
* Contains 8.25 units of Slow Release Nitrogen.	
<b>ACTIVE JNGREDIENT:</b>	
Atrazine: CAS(L912-24-9).....	00.92%
<b>INERT INGREDIENTS:</b> .....	<b>99.08%</b>
<b>TOTAL.....</b>	<b>100.00%</b>
This product contains 18.40lbs. of Atrazine per ton. This is equivalent to an actual application rate of 2 lb. active ingredient Atrazine per acre when used as directed.	

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
**NET WEIGHT 50 LB (22.68 kg)**

**General Precautions and Restrictions For Best Results**

- **Apply no more than 25lbs per 5,000 sq. ft.** per application, which is equivalent to 2 pound active ingredient atrazine per acre. A second application, (maximum of 2 per year) to **same area may give better results.**
- Do not apply in such a manner that will contact any person or pet whether directly, or through drift. Keep people and pets out of the area during application.
- Weeder Feeder must be watered into the turf immediately after application for safety, best weed control, and fertilizer benefit. “Entry Restrictions, Do not enter or allow others (including children or pets) to enter the treated areas (except those involved in the watering) until the watering-in is complete and the surface is dry”.
- Do not apply by hand. Avoid contact with hands or skin. Broadcast applications must NOT be made using hand-held devices, such as a belly grinder or handled rotary applicator. Such equipment may only be used for spot treatments.

WEEDS CONTROLLED OR SUPPRESSED		
Annual Bluegrass (Poa Annual)	Cutleaf Evening Primrose	Oxalix (Common and Yellow Woodsorrel)
Annual Sowthistle	Dichondra	Parsley-Pieri
Annual Clover	Dollarweed	Paspalums
Black Medic	Florida Belony (Rattlesnake Weed)	Red Sorrel
Blue Toadflax	Florida Pursley	Rescue Grass
Bull Paspalum	Fragrant Cudweed (Rabbit Tobacco)	Sandspur
Buckhorn Plantain	Henbit	Shepherdspurse
Bur Clover	Knotweed	Spurweed
Carolina Geranium (Cranesbill)	Lespedeza	Stinging Nettle
Chickweed (Common & Mouse-Ear)	Mat Lippia (Matchweed, Creeping, Charlie	Swinecrest
Dandelion	Moneywort	Venus Looking Glass
Common Purslane	Narrowleaf Vetch	Virginia Creeper
Grabgrass		Wild Mustard
Cudweed		

**AMOUNT TO USE**

Apply Sunniland Weeder Feeder at the rate of 25 lbs per 5,000 sq. ft. Water in after application. A second application in the same year, to the same area at the same rate will give longer weed control and required turfgrass response to the fertilizer.

**WHEN TO APPLY**

For best results apply before weeds come up. Established weeds are more difficult to control and may require an application in the spring and in the fall. Usually application made between January 1 and March 1 give best weed control; but applications may be made through June.

**IMPORTANT USE PRECAUTIONS**

DO NOT APPLY THIS PRODUCT ON OR OVER THE ROOT ZONE OF ORNAMENTAL TREES, SHRUBS, BEDDING PLANTS, FLOWERS OR GARDEN PLANTS OR INJURY TO THESE PLANTS MAY RESULT.

- Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result.
- Do not apply to newly seeded Zoysia, Centipede, and Carpetgrass lawns until after the new grass has been mowed at least twice.
- Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. May be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds.
- Do not plant any crop (flower or vegetable gardens ) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80 F or above water sufficiently to rinse particles off the grass blades.

SPREADER SETTINGS BROADCAST TYPE SPREADER SETTINGS	
FOR ONE PASS APPLICATION: For best results, apply in a circular path starting on the outside and working toward the center. A little overlap (1 to 2 inches) of product will ensure complete coverage for weed control and fertilizer benefit.	
5 LBS. PER 1,000 SQ. FT.	
Republic EZ Rotary	1 1/2
Scott's EasyGreen	25 1/4
Quaker Rotary	3 1/4
Ortho Rotary (6000)	1 1/2
EV-N Spred	12 1/2
True Temper	3 1/4
Precision Rotary	7 1/2
DROP SPREADER SETTINGS	
FOR ONE PASS APPLICATION: First apply 2 border strips along the edge of each end of the lawn. Then apply products along the length of the lawn, Overlapping slightly (1 to 2 inches) and shutting off the Spreader before turning as you reach the edge of the border strip.	
5 LBS. PER 1,000 SQ. FT.	
Ortho Drop	5
Republic EZ Drop	5
Scott's Drop (PF3)	5.5
Scott's Even Green	5.5