

6-0-15 9% Ca

Patented Amino Acid Sprayable Powder

Guaranteed Analysis

)%
)%
%

Derived from amino acids, calcium carbonate, calcium hydroxide, calcium oxide, calcium silicate, calcium sulfate

Amino Acid Profile			
Cystine	Alanine	Arginine	Aspartic Acid
0.6%	2.1%	2.7%	3.2%
Glutamic Acid	Glycine	Histidine	Isoleucine
4.7%	3.6%	0.5%	1.3%
Leucine	Lysine	Methionine	Phenylalanine
3.1%	0.7%	0.2%	2.2%
Proline	Serine	Threonine	Tyrosine
4.8%	5.5%	2.0%	0.6%
Valine	Amino Acid %:	N from Amino Acids:	
2.5%	40.3%	6.2%	

CAUTION: KEEP OUT OF REACH OF CHILDREN



WARNING

Causes skin irritation. Causes eye irritation.

PRECAUTIONARY STATEMENTS:

Wash skin thoroughly after handling. Wear protective gloves/protective clothing/eye and face protection.

IF ON SKIN: Wash with plenty of with soap and water. If skin irritation occurs: Get medical advice/attention.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists: Get medical attention/advice.

Take off contaminated clothing and wash it before reuse.

General Information

Pennamin® High-K powder contains 6% N, 15% K₂O and 9% Ca, as well as a blend of 17 different *L-amino acids*, 100% of which are free amino acids. Amino acids are a source of organic nitrogen, an essential nutrient for all plants and microorganisms, and are more easily broken down and digested.

Pennamin High-K is rapidly absorbed through the stomates and trans-located within the plant to correct calcium deficiencies and enhance general vigor.

General Recommendations

Pennamin products can be incorporated into liquid fertilizers or applied separately. Soil applications may be made either by sprinkler or band application.

Use a minimum of 5 gallons of water per pound of Pennamin.

Pennamin High-K can be used on turf, ornamentals, flowers, fruits, vegetables, tree and vine crops, row crops, field crops, trees, etc.

Application Rate:

Apply 1.0 to 5.0 pounds of **Pennamin High-K** per acre (0.4 to 2.0 ounces per 1.000 ft^2).

For trees, tree crops and vines crops dilute 0.5 - 3.0 pounds of **Pennamin High-K** in 125 gallons of water.

Timing of foliar sprays:

- Apply every 14 to 28 days as needed.
- Applications on turf and ornamentals can be made through out the growing season.

Note: Frequent applications at lower rates are generally more effective than fewer applications at higher rates.

Mixing Instructions

Fill tank with half the water, add **Pennamin High-K** while agitating solution, then add balance of water.

Compatibility

DO NOT mix with Phosphates

Do not mix with pesticides or other chemicals without a compatibility test. **Pennamin High-K** is compatible with most common foliar pesticides and nutrients; however, USER ASSUMES FULL RESPONSIBILITY to ensure compatibility when tank mixing with other products.

DO NOT STORE SOLUTION

Conditions of Sale

Seller warrants this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions and use. No one other than an officer of the seller is authorized to make any warranty, guarantee, or direction concerning this product. Because time, place, rate of application and other conditions of use are beyond the seller's control, the seller's liability from handling, storage, and use of this product is limited to replacement of the product or a refund of the purchase price.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm or by writing to LidoChem, Inc. at the address on this label and referring to the lot/batch # on this container.

40 lb. (18.1 kg.) @ 68°F



Guaranteed and Distributed By:

LidoChem, Inc.

www.pnfertilizers.com